


P. N. LEBEDEV PHYSICAL INSTITUTE OF  
THE RUSSIAN ACADEMY OF SCIENCES



## PRESENTATIONS

### of 19th International Symposium on Very High Energy Cosmic Ray Interactions – ISVHECRI 2016

#### Abstracts



The Symposium is carried out under the  
auspices of ***the International Union of Pure  
and Applied Physics (IUPAP)***

with the financial, material and technical support  
from:



- ***the Federal Agency for Scientific Organizations (FASO);***
- ***the P.N. Lebedev Physical Institute of the Russian Academy of Sciences (LPI RAS);***
- ***the Russian Foundation for Basic Research (RFBR).***

**22 – 27 August 2016**

Moscow, Russia

# **Model-stable universality of the air shower electromagnetic component: an approach to solving the mass composition problem**

R.I. Raikin<sup>1</sup>, T.L. Serebryakova<sup>1</sup>, A.A. Lagutin<sup>2</sup>, N.V. Volkov<sup>1</sup>

<sup>1</sup>*Altai State University, Barnaul, Russia*

<sup>2</sup>*Altai State University, Radiophysics and Theoretical Physics Department, Barnaul, Russia*

## **Abstract**

New results on the universal relation between radial scale factor and longitudinal age of the extensive air shower electron component are reported. We confirm that the scale factor of electron lateral distribution is potentially effective primary mass estimator, being evaluated from the surface detectors data with high duty cycle. When combining with the fluorescence detectors data on longitudinal shower age, it provides improved composition results both for average shower measurements and on event-by-event basis. Using CORSIKA simulations, we examine the sensitivity of conclusions to the hadronic interactions model. The enhancements of such a theoretically motivated tool for the unbiased cosmic ray composition deduction in a wide primary energy range from multi-hybrid EAS data of nearest future are discussed.