

P. N. LEBEDEV PHYSICAL INSTITUTE OF  
THE RUSSIAN ACADEMY OF SCIENCES



## PRESENTATIONS

### of 19th International Symposium on Very High Energy Cosmic Ray Interactions – ISVHECRI 2016

#### Abstracts



The Symposium is carried out under the  
auspices of ***the International Union of Pure  
and Applied Physics (IUPAP)***

with the financial, material and technical support  
from:



- ***the Federal Agency for Scientific Organizations (FASO);***
- ***the P.N. Lebedev Physical Institute of the Russian Academy of Sciences (LPI RAS);***
- ***the Russian Foundation for Basic Research (RFBR).***

**22 – 27 August 2016**

Moscow, Russia



# Average value of the cosmic ray injection spectrum at the Galactic sources

A.A. Lagutin<sup>1</sup>, N.V. Volkov<sup>1</sup>

<sup>1</sup>*Altai State University, Barnaul, Russia*

## Abstract

Experimental data and theoretical calculations show that the interstellar medium (ISM) is highly non-homogeneous on the scales of hundreds of parsecs. Filaments, shells, clouds are entities widely spread in the ISM. Spatial structure of these entities changes with time. The particles emitted by galactic sources, propagating in such a medium en route to the Solar system pass through the regions of the Galaxy, that have the different diffusion coefficients.

In this paper we analyze the effect of the variations of the ISM characteristics during the diffusion of the particles from the sources to the Solar system on the observed energy spectra of cosmic rays. Using Ginzburg-Syrovatskii normal diffusion model, we show that the different scenarios of the spatial irregularities distribution lead to diffusion models that demonstrate anomalous properties.

Our calculations show that the agreement of the new anomalous diffusion models with the experimental data is reached with the average value of the Galactic sources spectrum index  $p \sim (2.7-2.9)$ .

The authors acknowledge support from the Russian Foundation for Basic Research grant No. 16-02-01103.